



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2019559 C 2002/01/08

(11)(21) **2 019 559**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) Date de dépôt/Filing Date: 1990/06/21

(41) Mise à la disp. pub./Open to Public Insp.: 1990/12/22

(45) Date de délivrance/Issue Date: 2002/01/08

(30) Priorité/Priority: 1989/06/22 (P 39 20 358.1) DE.

(51) Cl.Int.⁵/Int.Cl.⁵ C12N 15/62, A61K 39/395, C12N 15/13, C07K 16/46

(72) Inventeurs/Inventors:
Steinstraßer, Axel, DE;
Kuhlmann, Ludwig, DE;
Bosslet, Klaus, DE;
Seemann, Gerhard, DE;
Hermentin, Peter, DE

(73) Propriétaire/Owner:
BEHRINGWERKE AKTIENGESELLSCHAFT, DE

(74) Agent: BERESKIN & PARR

(54) Titre : RECEPTEURS BISPECIFIQUES ET OLIGOSPECIFIQUES MONO- ET OLIGOVALENTS, PREPARATION ET UTILISATION

(54) Title: BISPECIFIC AND OLIGOSPECIFIC MONO- AND OLIGOVALENT RECEPTORS, THE PREPARATION AND USE THEREOF

(57) Abrégé/Abstract:

The invention relates to bispecific and oligospecific, mono- and oligovalent receptors which are prepared by gene manipulation by fusion of DNA coding for F(ab) fragments of antibodies of two or more different specificities by means of suitable linkers. In this connection, one specificity is preferably directed either against an epitope, which is located on the cell membrane or in the interstitium, of a tumor-associated antigen (TAA) or against an epitope in the tumor endothelium (TE), while the other specificities relate to highmolecular or low-molecular weight ligands and react, for example, with the Komplexons ethylenediaminetetraacetate and diethylenetriaminepentaacetate in Y90 complexed-form (EDTA-Y90 and DTPA-Y90 respectively). In a particularly preferred embodiment, the binding with the Komplexons takes place on the Komplexon receptor arm via fos-jun interaction (or else avidin-biotin interaction). Other preferred specificities have catalytic properties.



Abstract

The invention relates to bispecific and oligospecific, mono- and oligovalent receptors which are prepared by gene manipulation by fusion of DNA coding for F(ab) fragments of antibodies of two or more different specificities by means of suitable linkers. In this connection, one specificity is preferably directed either against an epitope, which is located on the cell membrane or in the interstitium, of a tumor-associated antigen (TAA) or against an epitope in the tumor endothelium (TE), while the other specificities relate to highmolecular or low-molecular weight ligands and react, for example, with the Komplexons ethylenediaminetetraacetate and diethylenetriaminepentaacetate in Y90 complexed-form (EDTA-Y90 and DTPA-Y90 respectively). In a particularly preferred embodiment, the binding with the Komplexons takes place on the Komplexon receptor arm via fos-jun interaction (or else avidin-biotin interaction). Other preferred specificities have catalytic properties.

- 1 -

Bispecific and oligospecific mono- and oligovalent
receptors, the preparation and use thereof

Background of the Invention

The invention relates to bispecific and oligospecific,
5 mono- and oligovalent receptors which are prepared by gene
manipulation by fusion of DNA coding for F(ab) fragments
of antibodies of two or more different specificities by
means of suitable linkers. In this connection, one
specificity is preferably directed either against an
10 epitope of a tumor-associated antigen (TAA), which is
located on the cell membrane or in the interstitium or
against an epitope in the tumor endothelium (TE), while
the other specificities relate to high-molecular or low-
molecular weight ligands and react, for example, with the
15 chelates ethylenediaminetetraacetate and
diethylenetriaminepentaacetate in Y90 complexed form
(EDTA-Y90 and DTPA-Y90 respectively). In a particularly
preferred embodiment, the binding with the chelates takes
place on the chelate receptor arm via fos-jun interaction
20 (or else avidin-biotin interaction). Other preferred
specificities have catalytic properties, or bind to other
TAA's on the same tumour cell or to receptors on lymphoid
cells.

Bispecific antibodies have to date been prepared by the
25 following methods:

- chemical coupling of antibodies of diverse
specificity via heterobifunctional linkers (H.
Paulus, Behring Inst. Mitt. 78, (1985), 118-132)
- fusion of hybrids which are already available and
30 secrete various monoclonal antibodies (MAb), and
isolation of the bispecific monovalent portion (U.S.
Staerz and X.J. Bevan, Proc. Natl. Acad. Sci. USA 83,
(1986) 1453-1457).

- 2 -

- transfection of the light and heavy chain genes of two different MAbs (4 genes) into murine myeloma Pells or other eukaryotic expression systems and isolation of the bispecific monovalent portion (U. Zimmermann, Rev. Physio. Biochem. Pharmacol. 105 (1986), 176-260; J. van Dijk et al., Int. J. Cancer 43, (1989) 1 344-349).

Bispecific antibodies of this type are employed for the therapy and diagnosis of malignant tumors. The principle of the process comprises, in the first step, achieving saturation of the epitopes which are recognized by one of the two specificities on the target cells, by injection of the bispecific macromolecule over prolonged periods and with high doses.

- 15 In the second step, which comprises interruption of the treatment for several days, there is autoelimination of the non-specifically adsorbed bispecific antibody from the nontarget tissues. This autoelimination can be speeded up by injection of an anti-idiotypic antibody which is coupled to sugar residues, preferably galactose, and is directed against the anti-tumor arm of the bispecific receptor.

The third step in the process comprises i.v. injection of a radiolabeled, hydrophilic low-molecular weight ligand which does not accumulate in cells, has a short residence time in the body, has high complexing constants for beta- and gamma-emitters such as ^{90}Y , ^{186}Re , ^{188}Re , ^{189}Re , $^{99\text{m}}\text{Tc}$ or ^{111}In , and binds to the second specificity of the bispecific receptor with high affinity. This step results in a concentration of the radioactive ligand, associated with prolonged retention on the target tissue, which results in selective destruction of the target tissue and makes possible diagnosis of metastases, for example.

- 3 -

Summary of the Invention

The present invention relates to bispecific or oligospecific receptors produced by gene manipulation. In particular, the present invention provides a bispecific or
5 oligospecific mono- or oligovalent receptor produced by gene manipulation comprising: (a) a V_H and C_{H1} region of a first antibody, which is directed against an tumour associated antigen; (b) a V_H and C_{H1} region of a second antibody, which is directed against a chelate, wherein the
10 chelate binding takes place on a receptor chelate arm via fos-jun interaction; (c) a polypeptide spacer that links C_{H1} region of the first antibody to the V_H region of the second antibody without impeding association with light chains of said antibodies or antigen binding; and (d) the
15 light chains of the first and second antibodies.

In a preferred embodiment, the receptor of the present invention further comprises: (e) a V_H and C_{H1} region of a third antibody, wherein the binding specificity of the third antibody is different from the antigen binding
20 specificity of the first and second antibody; (f) a polypeptide spacer that links the third antibody to the first or second antibody; and (g) the light chain of the third antibody. In a specific embodiment, the binding specificity of the third antibody is directed against DTPA
25 or EDTA.

The present invention also relates to a process of preparing the receptor of the invention comprising the steps of: (a) preparing an expression vector encoding a receptor of the invention; (b) preparing an expression
30 vector expressing the light chains of a receptor of the invention; (c) transforming cells with the expression vector from (a) and the expression vector of (b); and (d) growing cells from (c) and isolating the receptor from the cells.

- 4 -

The receptors of the invention are useful as pharmaceuticals.

Brief Description of the Drawings

The invention will now be described in relation to the
5 drawings in which:

- Figure 1 is a schematic of a receptor of the invention.
Figure 2 is a schematic of a receptor of the invention.
Figure 3 is a schematic of a receptor of the invention.
Figure 4 is a schematic of a receptor of the invention.
10 Figure 5 is a schematic of the plasmid IgG3C F(ab')₂1H.
Figure 6 is a schematic of the plasmid M.
Figure 7 is a schematic of the plasmid V_Ha/V_Hb PcR.
Figure 8 is a schematic of the plasmid PEV_HaΔC.
Figure 9 is a schematic showing the preparation of plasmid
15 L.
Figure 10 is a schematic showing the preparation of
plasmid X.
Figure 11 is a schematic showing the preparation of
plasmid Z.
20 Figure 12 is a schematic showing the preparation of
plasmid Y.
Figure 13 is a schematic showing the preparation of
plasmid PEVT.
Figure 14 is a schematic of the plasmid D.
25 Figure 15 is a schematic of the plasmid E.
Figure 16 is a schematic showing the preparation of
plasmid F.
Figure 17 is a schematic showing the preparation of the
plasmid G (M13mp18).
30 Figure 18 is a schematic showing the preparation of
plasmid G'.
Figure 19 is a schematic showing the preparation of
plasmid H.
Figure 20 is a schematic showing the preparation of
35 plasmid I.

- 5 -

Figure 21 is a schematic showing the preparation of plasmid K.

Figure 22 shows the nucleic acid and amino acid sequence of peptide linkers.

- 5 Figure 23 shows the nucleic acid sequence and amino acid sequence of the V_H and V_K regions of MAb a.

Figure 24 shows the nucleic acid sequence and amino acid sequence of the V_H and V_K regions of MAb b.

- 10 Figure 25 shows the nucleic acid sequence and amino acid sequence of the V_H and V_K regions of MAb c.

Figure 26 shows the nucleic acid sequence and amino acid sequence of the V_H and V_K regions of MAb d.

Figure 27 shows the nucleic acid sequence of mutagenic oligonucleotides.

- 15 Figure 28 shows the nucleic acid and amino acid sequence of the mutated hinge exon.

Figure 29 shows the nucleic acid sequence of the JunI and JunII oligonucleotides.

Detailed Description of the Invention

- 20 The invention now provides bispecific and oligospecific receptors which have, depending on requirements, mono- or oligovalent binding sites to the particular epitopes and are produced by gene manipulation by means of suitable linkers. This entails the gene fragments which code for the V_H and C_{H1} sections of antibodies a and b being
- 25 linked, by means of suitable synthetic oligonucleotides as depicted by way of example in Figure 22, in such a manner that the N-terminus of the V_H domain of MAb b is covalently bonded via a polypeptide spacer to the C terminus of the C_{H1} domain of Mab a (Fig. 1). The V_HaC_{H1a} -polypeptide spacer - V_HbC_{H1b} gene construct is transfected
- 30 together with the genes for the light chains of antibodies a and b into eukaryotic cells (for example mouse myeloma cells). The C_{H1a} , C_{H1b} , C_{k1a} and C_{k1b} domains are modified in
- 35 such a way that opposite charges meet at the areas of

- 6 -

contact of the constant domains ($C_{H1a}(+)C_{k1a}(-)$; $C_{H1b}(-)C_{k1b}(+)$) (+ = positive, - = negative) or opposing areas of contact are in each case hydrophobic or in each case hydrophilic. This means that the transfectomas
5 preferentially express hybrid molecules which have the correct pairings of heavy and light chains (Fig. 2).

Here, antibody a is a representative example of an antitumor antibody, and antibody b represents an antibody
10 against a low- molecular weight ligand, preferably the chelates DTPA-Y90 or EDTA- Y90.

Bi- or oligospecific receptor accordingly means a genetically engineered construction of V_H and C_{H1} domains of antibodies of diverse specificity via suitable linkers so that the required mobility for association with the
15 corresponding light chains is present and antigen binding is not impeded.

The antigen-binding sites are called valencies or binding sites. A bispecific monovalent receptor thus has one antigen-binding site in each case when there are two
20 specificities. Consequently, a bispecific trivalent receptor has one antigen-binding site for one specificity and two antigen-binding sites for the other.

Bispecific receptors which are bivalent for the tumor antigen (Mab a) and monovalent for EDTA-Y90 (Mab b) are
25 prepared by linking the heavy chain gene construct described above, by means of the abovementioned oligonucleotide linker, to the gene section which codes for the V_H and C_{H1} domains of Mab a (Fig. 3) so that the C- terminal end of the C_{H1} domain of Mab b is connected to
30 the N- terminal end of the V_H domain of Mab a by a polypeptide spacer. These gene constructs are transfected together with the genes for the light chains belonging to Mab a and b into eukaryotic cells (for example myeloma

- 7 -

cells). The C_{H1} and C_{κ} domains are provided, as described above, with complementary charges or areas of contact which are in each case hydrophobic or hydrophilic. The transfectomas preferentially express fusion molecules which comprise two F(ab) fragments of MAb a and one F(ab) fragment of MAb b (Fig. 4). The mobility of the peptide linkers makes possible the alignment of the two F(ab) arms of MAb a towards the tumor cell with, at the same time, alignment of the F(ab) arm of MAb b towards the intercellular space. It is possible correspondingly to add further binding sites of identical or different specificity. Moreover, it is possible to combine freely the sequence of specificities in the constructs.

Consequently, the invention relates to bispecific or oligospecific, mono- or oligovalent receptors which have both specificity for an epitope located on the cell membrane or in the interstitium, for example TAA or TE, and specificity for a low- or high-molecular weight ligand which is distributed exclusively in the extracellular space. In this connection, one specificity is preferably formed by the tumor-specific antibodies, whereas the other specificity is preferably directed against DTPA-Y90 or EDTA-Y90. In a particularly preferred embodiment, the binding takes place with chelates on the chelate receptor arm via fos-jun interaction (see Example 5). Another preferred variant of the invention comprises the incorporation of specificities with catalytic activity. Moreover, there is no restriction on the choice of the sequence of the specificities or binding valencies, as is shown by way of example in Fig. 4 for three valencies with two specificities.

Particularly preferred constructs according to the invention are those which contain a V gene of Figures 23, 24, 25 and/or 26. Antibodies with these sequences and their properties. Moreover, the complementarity

- 8 -

determining regions (CDRs) can be identified by the method of Kabat and Wu (Sequences of Proteins of Immunological Interest, US Dept. of Health and Human Services, US Government Printing office (1987)). Likewise preferred are
5 constructs which contain specificities against the epitopes defined by the monoclonal antibodies described above.

The invention additionally relates to genetic engineering processes for the preparation of the constructs described
10 above, and to a use of the abovementioned constructs for preparing pharmaceuticals for controlling and diagnosing target cells. This entails, in a first step, saturation of the relevant epitopes on target cells after injection of the constructs and, in a subsequent interval, elimination
15 of non-specifically adsorbed or unbound constructs. The step following this comprises injection and subsequent specific binding of a low- or high-molecular weight ligand which does not accumulate in cells and is intrinsically cytotoxic or is "activated" to cytotoxicity by
20 extracorporeal influences where appropriate in another step. Examples of processes of this type are enzymatic activation, activation by microwave irradiation of a prodrug or activation by laser light.

The invention is furthermore contained in the examples and
25 the patent claims.

Example 1: Preparation of an anti-DTPA-Y90 or EDTA-Y90 MAb

Isothiocyanatobenzyl-DTPA (formula 2) was covalently coupled as hapten onto human serum albumin (HSA as carrier) with a degree of derivatization of 19 benzyl-DTPA
30 molecules per HSA molecule by the method described in (N.W. Brechbiel et al., Inorganic Chemistry 25, (1986) 2772-2781). 20 µg of this hapten-carrier complex, into which cold Y had been complexed, were injected s.c. on day

- 9 -

0 with Freund's adjuvant, on day 7 and 14 with incomplete Freund's adjuvant and on day 21 with PBS into Balb/c mice. on day 24, the spleens of the mice with the highest anti-DTPA antibody titers were fused with the SP2/0-Ag14 myeloma cell line (Shulman et al., Nature 276, (1978) 5 269). The resulting hybridomas were tested in a DTPA-specific ELISA for the production of high-affinity MABs against DTPA and EDTA. The ELISA comprised a solid phase which was loaded with a solution containing HSA-benzyl-10 DTPA-Y. The supernatant containing the MAB to be tested was preincubated with free chelate or its metal ion complexes, and its binding to the specific solid phase was measured. An enzyme amplification system which is coupled to an anti-mouse immunoglobulin antibody was used for this 15 purpose. The details of these methods are described in Annex 1a and 1b.

MABs which have the properties described in Annex 1e were obtained using this assay system.

20 In contrast to many other anti-DTPA/EDTA MABs, these MABs do not bind to normal human tissue, as was found using the APAAP technique (Cordell et al., J. Histochem. Cytochem. 32: 219, 1984) on cryopreserved tissues. It is therefore possible to use these MABs in vivo for diagnosis and therapy.

25 The competitors employed were the chelates DTPA and EDTA in non-complexed and in complexed form (Annex 1c).

In addition, the structurally related compounds transaconitic acid and 1,2-diaminoethane were used as inhibitors (see Annex 1e). MAB BW 2050/174 is particularly 30 suitable for in vivo use, showing preferential binding to EDTA-Y, in contrast to all the other MABs (see Annex 1e, low competitor excess for EDTA-Y (100 x) higher excess for other EDTA Komplexons). The hybrid 2050/174 was therefore

- 10 -

stabilized and used for developing the EDTA-Y arm in the bispecific receptor.

Example 2: Preparation and expression of a V_{H1a} C_{H1} -linker- V_{H1b} C_{H1} gene construct

- 5 The techniques used here were taken, unless indicated otherwise, from Molecular Cloning, A Laboratory Manual; Sambrook, Fritsch, Maniatis; Cold Spring Harbor Laboratory, 1982 (pp. 11-44, 51-127, 133-134, 141, 146, 150-167, 170, 188-193, 197-199, 248-255, 270-294, 310-328, 10 364-401, 437-506) and from Molecular Cloning, A Laboratory Manual, Second Edition; Sambrook, Fritsch, Maniatis; Cold Spring Harbor Laboratory Press, 1989, (pp. 16.2-16.22, 16.30-16.40, 16.54-16.55).

- A human IgG_3 C gene was isolated from a human gene bank in 15 EMBL3 phages (A.M. Frischau et al., J. Mol. Biol. 170, 827-842 (1983) and G.H.A. Seemann et al., The EMBO Journal 5 (1986), 547-552).

- Constructions which contain, on the one hand, only the C_{H1} exon and a hinge exon (Fig. 5) and, on the other hand, the 20 C_{H1} exon and the 3'NT region of an HLA B27 gene (Fig. 6, fragment M in plasmid M) were prepared from this IgG_3 C gene as described in German Patent Application P 38 25 615.0.

- The V_{H1a} and V_{H1b} genes were amplified from mRNA of the 25 hybrid clones a and b as described by Orlandi et al. (Proc. Natl. Acad. Sci. USA 86, (1989), 3833-3837) and cloned in an M13 vector (V_{H1a} PCR and V_{H1b} PCR) (Fig. 7). The V_{H1a} gene was cloned as the HindIII-BamHI fragment into the eukaryotic expression vector pEV_H (Simon et al., Nucl.

- 11 -

Acids. Res. 16, (1988), 354) (Fig. 8). The result is the plasmid pEV_H a C.

The human IgG C gene subclone with the C_H1 and with one hinge exon (Fig. 5) contains a PstI cleavage site between
5 C_H1 exon and hinge exon. The V_H genes contain a PstI cleavage site at the 5' end. The linker oligonucleotide is designed such that it overlaps at the 5' end with the region of the PstI cleavage site on the C_H1 + 1H subfragment of the IgG C gene and at the 3' end with the
10 PstI cleavage site of the V_Hb gene. The linker oligonucleotide is cloned by means of its PstI cleavage sites into the PstI cleavage site of a PUC 18 plasmid (Fig. 9). The result is the plasmid clone L.

The plasmid with the IgG₃ C gene subfragment with C_H1 exon
15 and with a hinge exon is cleaved with PstI and BamHI and ligated to the V_Hb gene fragment cut out of V_Hb PCR as PstI-BamHI fragment (Fig. 10). The result is the plasmid X.

The plasmid X is cleaved with PstI and ligated to the
20 linker fragment which has been cut out of the plasmid L likewise with PstI (Fig. 11). Nucleic acid sequence analysis is used to identify the clone Z in which the linker is cloned in correct orientation between C_H1 and V_Hb without disturbing the intron/exon junction between
25 intron 3 and linker exon and without disturbing the reading frame at the junction between linker and V_Hb gene.

The plasmid pEVa C is cleaved with BamHI and ligated to the f ragment M cut out of the plasmid M with BamHI. Restriction analysis is used to identify the clone Y which
30 contains the fragment M in the correct orientation (Fig. 12).

- 12 -

The plasmid Y is partially cleaved with BamHI and ligated to the fragment (C_H1-linker-V_Hb) cut out of the plasmid X by HindIII and BamHI cleavage, after all the ends have been filled in. Nucleotide sequence analysis and
5 restriction mapping are used to identify the plasmid clone PEVT which contains the fusion gene V_Ha C_H1-linker-V_Hb C_H1 with the correct orientation of all the exons (Fig. 13).

The plasmid PEVT is transfected together with plasmids which harbor the genes for the light chains of antibodies
10 a and b into suitable eukaryotic expression cells in order to express the antibody a F(ab) antibody b F(ab) fusion protein.

Example 3: Transfection of the light and heavy chain genes of two different MAb (4 genes)

15 The genes cloned into vectors were transfected by electroporation after linearization of the vectors into X63Ag8.653 myeloma cells (H. Stopper et al., Biochem. Biophys. Acta 900 (1987), 38-44). The transfectomas which grew in selective media were tested for the production of
20 bispecific monovalent MAb in a specific RIA. This RIA comprised TAA adsorbed on a solid phase, onto which, after blockade of the non-specific sites by casein, the transfectoma supernatants to be assayed were placed. After DTPA or EDTA which were complexed with ⁹⁰Y or ^{99m}Tc had
25 been added and the excess had been washed away, it was possible to detect those transfectomas which secreted bispecific monovalent anti-TAA x anti-EDTA MAb by an increased radioactive signal on the solid phase.

Transfectoma 9 was stabilized by limited dilution cloning
30 and expanded in cell culture. Cell culture supernatants were concentrated 10 x, the MAb fraction was purified by

- 13 -

protein A chromatography (P.L. Ey et al., Immunochemistry 15, (1978), 429), and the fraction containing the bispecific monovalent MAb was purified by anion exchange chromatography (J. Van Dijk et al., Int. J. Cancer 43, 5 (1989), 344-349).

Example 4: Biological effectivity

Purified protein containing the bispecific monovalent MAb (BW 431/26 x BW 2050/174) was injected i.v. in 500 µg doses on days 0, 3, 5, 8, 10 and 12 into nude mice carrying human tumor xenografts (CoCa 4). 50 µCi of EDTA-Y90 were injected i.v. into each of the animals of day 27-30. A 2nd group of animals received on the same days 500 µg of MAb BW 431/26 in place of the bispecific MAb, and the EDTA-Y90 injections as described above.

15 A third tumor-carrying group received injections of PBS (as tumor- growth control) in place of the MAb and EDTA-Y90. Tumor growth was followed for 6 weeks. Injection of EDTA-Y90 resulted in significant inhibition of tumor growth in the group which received the bispecific monovalent MAb, whereas the animals injected with MAb BW 20 431/26 and treated with EDTA-Y90 showed no inhibition of tumor growth, compared with animals which received only PBS.

25 These data indicate the selective tumor-therapeutic efficacy of the bispecific monovalent MAb in combination with EDTA-Y90 as toxic principle.

30 Even more favorable tumor-therapeutic effects are obtained by the oligovalent/bispecific or oligospecific receptors because they remain longer on the tumor, because of the bivalent binding to TAA, and thus the ligand is likewise retained on the tumor for longer and in higher concentrations.

- 14 -

Example 5: Optimization of the biological effectivity of bi- or oligospecific macromolecules by increasing the avidity of the anti- chelate arm.

A crucial factor which influences the efficient attachment of the hydrophilic chelate undergoing extracellular distribution to the anti- chelate arm of the oligospecific macromolecule is the avidity of this arm for the chelate. The avidities of monoclonal antibodies for their corresponding epitopes are in the range 10^5 - 10^{11} l/mol. Since these binding strengths are possibly insufficient to localize on the tumor the mass of chelate necessary for efficient radioimmunotherapy, in the following example the extremely strong interaction of the fos-leucine-zipper peptide (fos-peptide) with the Junleucine-zipper peptide (Jun- peptide) (Erin K. O'Shera et al., Science, 245, 1989) was used to immobilize the chelate as firmly as possible on the anti- chelate arm. It is necessary, in order to be able to utilize this strong fos-jun interaction, preferably to link the fospeptide covalently to the chelate (DTPA). It is possible for this purpose to react in a first step isothiocyanatobenzyl-DTPA with hydrazine (or a diaminoalkane). The DTPA-benzylthiocarbazide produced in this way can be reacted, in a 2nd step, with N- (gamma-maleimidobutyryloxy) succinimide or an analog to give DTPA-benzyl (gamma-maleimidobutyryl) thiocarbazide. Then, in a 3rd step, this compound is linked to the fos-peptide which has been extended by glycine-glycine-cysteine (Figure 1) via the free SH group of the amino-terminal cysteine. The fos-peptide-DTPA conjugate produced in this way is complexed in a 4th step with yttrium chloride. The fospeptide-DTPA-Y conjugate complex produced in this way can then be used for in vivo addition onto the Jun-peptide arm of the bi- or oligospecific macromolecule. The synthesis of the example outlined above is described in detail hereinafter:

- 15 -

A) Preparation of the fOB-EDTA-Y conjugate complexStep 1:Synthesis of EDTA-benzylthiocarbazide

5 Isothiocyanatobenzyl-EDTA (SCN-Bn-EDTA) (30 mg, 54 μ mol)
was stirred in 10% (v/v) aqueous hydrazine for 1 h. The
solvent was then removed under high vacuum, and the
residue was dried over phosphorus pentoxide under high
vacuum and finally freeze-dried. The product was
neutralized with DOWEX™ WX 2 (H⁺ form) and again freeze-
10 dried (yield 28 mg).

Step 2:Synthesis of EDTA-benzyl(gamma-maleimidobutyryl)thio-
carbazide

The EDTA-benzylthiocarbazide prepared in step 1 (20 mg; 34
15 μ mol) and N-(gamma-maleimidobutyryloxy)succinimide (8 mg,
29 μ mol = 0.9 equiv.) were stirred in anhydrous
dimethylformamide for 1 h. The mixture was then evaporated
to dryness, and the residue was dried under high vacuum.

Step 3:20 Coupling of the EDTA-benzyl(gamma-maleimidobutyryl)thio-
carbazide to the amino-terminal cysteine in the fos-
peptide

A solution of the fos-peptide (4.8 mg, 1 μ mol) (see step
3.1) in phosphate-buffered saline (2 ml) was mixed with a
25 suspension of the product mixture obtained as in step 2 (4
mg) in dimethylformamide (400 μ l) and incubated at room
temperature for 1 h. The reaction mixture was then
subjected to gel filtration on a Sephadex™ G15 column in
phosphate-buffered saline. The protein- containing eluate
30 was collected and preserved at -30°C (yield 4.2 mg).

- 16 -

Step 3. 1:

Amino acid sequence of the fos-peptide (| |) with N-terminal GGC extension.

Ac-CGGyLTDTLQAETDQLEDKKSALQTEIANLLKEKEKLEFILAAY

- 5 The letters represent the following amino acids: A = alanine, C = cysteine, D = aspartic acid, E = glutamic acid, G = glycine, I = isoleucine, K = lysine, L = leucine, M = methionine, N = asparagine, Q = glutamine, R = arginine, S = serine, T = threonine, V = valine, Y =
10 tyrosine.

- The oligopeptide was synthesized using an automatic peptide synthesizer (Applied Biosystems Model 430A) by the Merrifield solid-phase method (Stewart and Young, Solid Phase Synthesis, Pierce Chemical Company, 2nd edition,
15 Rockford Ill.) with the tert- butyloxycarbonyl protective group. The oligopeptides were cleaved off the phenyl acetamidomethyl -polystyrene support. After elimination of the protective groups (Tom et al., 1983, J. Am. Chem. Soc. 105, 6442-6455) the oligopeptides were purified by
20 reversed phase chromatography (PepRPC™ column, Pharmacia) as described by Rivier et al. (J. Chromatography 288, 303-328, 1984).

Step 4:

- Preparation of a fos-Deptide-EDTA-yttrium chelate with a
25 fos- peptide-EDTA conjugate prepared as in step 3

- The fos-peptide-EDTA conjugate prepared as in step 3 (4.2 mg) was dialysed against isotonic saline/0.1 M sodium citrate, pH 7.0, in a dialysis tube with an exclusion limit of m.w. 1,000 (Spectrum), and was mixed with 6 mg of
30 yttrium chloride which were dissolved in 3 ml of isotonic saline/0.1 M sodium citrate, pH 7.0. After 1 h, back-dialysis against phosphate-buffered saline was carried out, and the chelate solution was preserved at -30°C. The fos- peptide-EDTA-Yttrium chelate described in the above

- 17 -

example is then used as ligand in order to bind with high avidity to the jun- peptide arm of the bi- or oligospecific macromolecule. The construction of a bispecific macromolecule particularly suitable for this interaction is described in the following example.

B) Construction of the MAb-jun fusion molecule

The techniques used here were taken, unless otherwise indicated, from Maniatis et al. (Laboratory Manual EMBL (1982), Heidelberg), and Sambrook (Molecular Cloning: A Laboratory Manual).

Step 1:

A human IgG3 C gene was isolated from a human gene bank in EMBL 3 phages (A.M. Frischauf et al., J. Mol. Biol. 170, 827-842, 1983 and G.H.A. Seemann et al., The EMBO Journal 5, 547-552, 1986). A construction (D) which contains only the CH1 exon and the first hinge exon of the IgG3 C gene (Fig. 14) was prepared from this IgG3 C gene as described in German Patent Application P 3825615.0, published February 1, 1990.

A human HLA B27k gene was isolated from the same gene bank as likewise described in German Patent Application P 3825615.0, published February 1, 1990. A construct (E) which contains only the C3 exon and the 3' NT region of the HLA B27k gene (Fig. 15) was prepared from this HLA B27k gene.

Step 2:

The C1 exon and the 3' NT region of the HLA B27k gene were cut out of the plasmid E with XbaI, and the fragment was isolated and cloned into the XbaI cleavage site of the construct D. Restriction analysis and nucleic acid sequence analysis were used to identify the clone F which contains the C3 exon and the 3' NT region of the HLA B27k

- 18 -

gene in the correct 5'-3' orientation 3' from the IgG3 C gene fragment (Fig. 16).

Step 3:

5 The insert of the clone F is cut out of the plasmid using the endonucleases HindIII and EcoRI and cloned between the HindIII and EcoRI cleavage sites of an M13mp18 double-stranded (DS) phage. The phage clone G which contains the antibody/HLA fusion gene fragment is isolated (Fig. 17).

Step 4:

10 Uracil single strands are prepared from the phage clone G by the method of T.A. Kunkel, 1985, Proc. Natl. Acad. Science, U.S.A., 82, 488-492. The single-stranded phages were hybridized with the mutagenic oligonucleotides 1 and 2 (Figure 27), and the gaps between the oligonucleotides
15 were closed with Klenow DNA polymerase and T4 ligase. After transformation into E. coli, restriction analysis and nucleic acid sequence analysis were used to identify a phage clone (G) in which the SstI restriction cleavage site at the 5' end of the hinge exon had been deleted. At
20 the same time, a SstI and a SphI restriction cleavage site were introduced at the 3' end of the hinge exon (Fig. 18). To delete the SstI cleavage site, the third base of the 2nd codon of the hinge exon was converted from C into G, and to introduce the SstI and SphI cleavage sites, the
25 bases 5'GAGCTCGGGGCA3' were introduced between the 15th and 16th codon of the hinge exon (Figure 28).

Step 5:

Double-stranded DNA of the phage clone G' is cleaved completely with SphI and partially with SstI. The
30 synthetic oligonucleotides Jun I and Jun II (Figure 29) are combined to give a double-stranded DNA fragment which contains at each of its ends a cut SphI and SstI

- 19 -

restriction cleavage site and codes for a peptide which contains the Jun leucine zipper (O'Shea et al., Science, 245, 646- 648, 1989).

5 The double-stranded DNA fragment is cloned into the SstI and SphI restriction cleavage sites of the F' phage clone, and the phage clone H which contains a gene construct in which the sequence for the Jun zipper peptide is inserted in the hinge exon is identified (Fig. 19).

Step 6:

10 The insert of the DS phage H was cut out with the restriction endonucleases HindIII and EcoRI, the ends were filled in with T4 polymerase and cloned into an SmaIcleaved KsF vector (Stratagene, 11099 North Torrey Pines Road, La Jolla California 92037). Plasmid clone I
15 which contains the antibody/ Jun/HLA fusion gene in the orientation (Fig. 20) in which it is flanked on both sides by a BamHI cleavage site was identified.

Step 7:

The antibody/Jun/HLA fusion gene was cut out of the KS
20 clone I with BamHI and cloned into the expression plasmid pABStop (Behringwerke AG) which contains a specific functional immunoglobulin V gene. The specific V gene was obtained as described in Patent Application P 3909799.4. The expression plasmid K which contains the
25 antibody/Jun/HLA fusion gene construct in the correct orientation downstream of the V. gene was identified (Fig. 21).

Cotransformation of the plasmid K with a plasmid which contains the gene for the light chain of the specific MAb,
30 and a plasmid which carries a resistance gene, leads to expression of a specific antibody F(ab')₂ fragment which contains in the hinge region two Jun zipper peptides, with

- 20 -

the Jun zipper peptide being modified in such a way that there is no longer any homodimer (Jun/Jun) formation.

Example 6:

5 Optimization of the amount of bi- or oligospecific
receptor on the tumor and minimization thereof in the
blood and normal tissues

Scientific investigations by others have shown that penetration of solid tumors by macromolecules > 50 kDa takes place slowly, and usually only the edge region or a
10 few areas in the tumor are reached. These investigations are based on experiments which comprise a single injection of small amounts of macromolecules. In contrast thereto, we have found that substantial penetration of the entire tumor mass in nude mouse xenografts is possible by a
15 repetitive i.v. injection of large amounts of bi- or oligospecific receptors (10 x 250 µg receptor/mouse for 10 days). Furthermore, because of their specific binding to TAA, the bi- or oligospecific receptors remain attached for long periods (> 20 days) in large amounts on the tumor
20 cell membrane and in the tumor interstitium. These results were obtained using the indirect alkaline phosphatase technique on cryopreserved thin sections of human colonic and pancreatic tumor xenografts.

During this time (after only 10 days) the bi- or
25 oligospecific receptor molecules had already been eliminated from the TAA- negative normal tissues and the blood by degradation and excretion. In order to shorten this elimination period, an anti- idiotypic MAb (anti Id) which reacts only with the anti-TAA arm of unbound bi- or
30 oligospecific receptor molecules was injected i.v. (1 x 50 µg of anti Id) 24 hours after completion of the ten injections of bi- or oligospecific receptors. This single injection speeded up the elimination of the unbound bi- or

- 21 -

oligospecific receptor molecules from the blood and increased the metabolism rate in liver and spleen.

It is possible on the basis of this manipulation to inject the chelate (EDTA-Y90) only 4 days after completion of the phase of penetration and binding of the bi- or oligospecific receptor. The following treatment regimen (for nude mice) is derived from these investigations:

- a) day 1-10, i.v. injection of 1 x 250 µg of bi- or oligospecific receptor each time
- 10 b) day 11, i.v. injection of 1 x 50 µg of anti Id
- c) day 14, i.v. injection of a therapeutic dose of EDTA-Y90.

On the basis of comparative immunoscintigraphic data in nude mice and tumor patients, this regimen ought to be suitable for tumor therapy in humans too. However, the amounts to be injected in the human system are in a different range of magnitude. 10 x 5-10 g of bispecific receptor, 1 x 1 g of anti Id. Injection of the anti Id is not indispensable for therapy.

20 Annex 1a
Quantitative inhibition ELISA for MAbS by DTPA or EDTA complexes

Material: divisible 96-well polystyrene microtiter plates (U shape) type B, from Nunc, No. 4-60445

- 25 1) 50 µl of Y-benzyl-DTPA-HSA 19 conjugate with a concentration of 1 µg of conjugate per ml of PBS, pH 7.2, are pipetted into each well and incubated at room temperature (RT) overnight.
- 2) The supernatant is removed by aspiration and washed 3 x with 0.05 M tris citrate buffer, pH 7.4, (wash
- 30

- 22 -

solution 1); (1 x wash = introduce 250 µl of wash solution per well, leave to stand for 2 min and remove by aspiration).

- 5 3) If the microtiter plate is not required immediately, it is left to stand (with the opening underneath) on cellulose at RT overnight. The plate is then sealed in films with drying cartridges (from Gaplast, Postfach 529, 8100 Garmisch-Partenkirchen). The plates can be kept at +4°C for at least 8 weeks under these conditions.
- 10 4) 250 µl of blocking solution are applied to each well and incubated at 37°C for 30 min.
- 15 5) Preincubation of the diluted hybridoma, supernatant with the competitor is carried out during the blocking (see Annex 2).
- 20 6) 50 µl of the appropriately prediluted and preincubated hybridoma supernatants to be tested are applied to each well and incubated at RT for 30 min.
- 20 7) Washing 3 x with wash solution 2 is subsequently carried out.
- 25 8) Subsequently 50 µl of goat anti-mouse IgG, antibodies which are labeled with alkaline phosphatase and have been diluted 1:500 in blocking solution are applied to each well and incubated at RT for 30 min.
- 25 9) Then washing 3 x with wash solution for Enzygnost is carried out.
- 25 10) Subsequently 50 µl of 0.1 mM NADP are added.

- 23 -

- 11) Incubation at RT is then carried out for 30 min.
- 12) During the incubation with NADP, the amplification system is made up as follows:
per plate 2 parts of INT and 1 part of PBS, pH 7.21
5 are mixed and 1 part of diaphorase and 1 part of ADH are pipetted in.
- 13) 50 μ l of this system are placed in each well.
- 14) When there is a distinct change in color from transparent to red the reaction is stopped with 100
10 μ l of a 0.1 N H_2SO_4 solution per well.
- 15) The extinctions are measured at 492 nm in a TITERTEK^R MULTISCAN. 50 μ l of NADP with 50 μ l of solution and 1200 μ l of 0.1 N H_2SO_4 are employed as blank.

15	NADP	- Sigma order No. N-0505
	INT	- Sigma order No. 1-8377
	ADH	- Sigma order No. A-3263
	DIAPHORASE	- Sigma order No. D-2381
	Wash solution 2	- Behring, order No. OSEW96
20		contains Tween TM /PBS

Blocking solution:

PBS, pH 7.2, is made 3% strength in casein by adding casein and stirring for 30 minutes, and is adjusted to pH 7.4. Particles are then removed by centrifugation at 4,000
25 rpm for 10'.

Diluted goat anti-mouse IgG₁ antibodies labeled with alkaline phosphatase (from Southern Biotechnology Associates, Cat. No. 1080-04).

- 24 -

Preparation of 0.1 mM NADP:

Dissolve 7.65 mg of NADP in 100 ml of 20 mM tris, 0.1 mM MgSO_4 , pH 9.5; this solution can be stored at -20°C for several months.

- 5 Preparation of INT (P-iodonitrotetrazolium Violet):
Dissolve 2.5 mg/ml of 30% ethanol in an ultrasonic bath; always make up fresh.

Preparation of diaphorase:

- 10 1 mg of diaphorase/ml of PBS, pH 7.2, is stored in portions at -20°C .

Preparation of alcohol dehydrogenase:

0.5 mg of ADH/ml of PBS, pH 7.2, are stored in portions at -20°C .

Annex 1b

- 15 Preincubation of the hybridoma supernatant with the competitor

The mouse IgG concentration in hybridoma supernatants can be determined using commercially available quantitative ELISA systems and is state of the art.

- 20 On the basis of the ELISA concentration determination, the hybridoma supernatants are diluted to 1.25 $\mu\text{g}/\text{ml}$ in PBS without Ca^{++} and Mg^{++} .

Conversion from gram into mol:

- 150,000 g - 1 mol of MAb
25 1.25×10^{-6} g - x mol

$$1.25 \mu\text{g} = x = 8.33 \times 10^{-12} \text{ mol}$$

- 25 -

In order to have a 1 + 1 ratio of MAb and inhibitor, 10 μ l of inhibitor with a concentration of 8.33×10^{-12} mol/200 μ l, which is increased by a factor of 5, were added to 50 μ l of hybridoma supernatant with a concentration of 8.33×10^{-12} mol/ml.

The hybridoma supernatant is incubated with 100,000-fold, 50,000-fold, 10,000-fold, 5,000-fold, 1,000-fold and 100-fold excess of competitor at RT for 301. 50 μ l of this are pipetted into the ELISA (see Annex 1a, no. 6).

10 Annex 1c

Production of the DTPA and EDTA complexes

The complexing constant of DTPA or EDTA to the metal ions depicted in Figure 22 is extremely high so that complete saturation has to be expected on equimolar mixing of DTPA or EDTA with these metal ions. For this reason, the corresponding metal ions were incubated in a 3-fold molar excess with the DTPA or EDTA. As an example, 170 μ l of a 10 mM cadmium sulfate solution in double-distilled water (see Annex 1d) were incubated with 30 μ l of a 0.028 molar DTPA stock solution in double-distilled water at RT for 5'. Mixing 10 μ l of this competitor solution with the hybridoma supernatant leads to a 100,000-fold excess of competitor over the MAb contained in the hybridoma supernatant. Lower competitor to MAb ratios were achieved by diluting the competitor solution in the particular salt ion solution appropriately for the desired molar excess (see Annex 1b).

- 26 -

Annex IdSource and relevant physicochemical parameters of the metal ions employed

10 millimolar solutions of the following metal ions were
5 prepared in double-distilled water:

	Manganese chloride	MW 161.88	
	from Merck	No. 5934	Mn ion radius: 80 pm
	Cadmium sulfate	MW 256.5	
	Riedel de Haen	No. 31145	Cd ion radius: 97 pm
10	Zinc chloride	MW 136.28	
	from Merck	No. 8816	Zn ion radius: 74 pm
	Copper sulfate	MW 159.61	
	from Riedel de Haen	No. 31294	Cu ion radius: 96 pm
	Yttrium chloride	MW 303.36	
15	from Aldrich	No. 20,491-9	Y ion radius: 92 pm
	Lead(II) nitrate	MW 331.20	
	from Riedel de Haen	No. 31137	Pb ion radius: 120 pm

Annex 1eQuantitative assay of inhibition of MAb by DTPA and EDTA

20 Molar excess of competitor which leads to 50% inhibition
of the binding to the solid-phase antigen.

- 27 -

	MAb No.	DTPA-Y	DTPA	DTPA-Mn	DTPA-Cd	DTPA-Zn	DTPA-Cu
	2050/174	10^4	10^3	10^2	10^2	5×10^3	5×10^3
	2050/531	5×10^4	10^3	10^2	10^2	5×10^3	5×10^3
	2050/532	5×10^4	10^3	10^2	10^2	5×10^3	5×10^3
5	2050/534	5×10^4	10^3	10^2	10^2	5×10^3	5×10^3
	2050/535	10^4	10^2	10^2	10^2	10^3	10^3


	MAb No.	DTPA-Pb	1,2-Diamino-ethane	Trans-aconitic acid	EDTA-Y	EDTA	EDTA-Mn
10	2050/174	10^3	no inhibition Up to 10^5	no inhibition up to 10^5	10^2	10^3	10^3
	2050/531	5×10^3	"	"	10^3	10^3	10^3
	2050/532	5×10^3	"	"	10^2	10^3	10^3
15	2050/534	5×10^3	"	"	10^2	10^3	10^2
	2050/535	10^3	"	"	10^2	10^2	10^2

	MAb No.	EDTA-Cd	EDTA-Zn	EDTA-Cu	EDTA-Pb
	2050/174	10^3	10^3	10^3	5×10^3
	2050/531	10^3	10^3	10^2	10^5
20	2050/532	10^3	10^3	5×10^3	10^5
	2050/534	10^2	10^3	10^3	5×10^3
	2050/535	10^2	10^2	10^2	10^2


Formula 1

- 28 -

DTPA

Formula 2

Isothiocyanatobenzyl-DTPA



- 29 -

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE
PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A bispecific or oligospecific mono- or oligovalent
receptor produced by gene manipulation comprising:
 - 5 (a) a V_H and C_{H1} region of a first antibody, which
is directed against an tumour associated
antigen;
 - (b) a V_H and C_{H1} region of a second antibody, which
is directed against a chelate, wherein the
10 chelate binding takes place on a receptor
chelate arm via fos-jun interaction;
 - (c) a polypeptide spacer that links C_{H1} region of
the first antibody to the V_H region of the
second antibody without impeding association
15 with light chains of said antibodies or antigen
binding; and
 - (d) the light chains of the first and second
antibodies.
2. A receptor as claimed in claim 1, wherein at least
20 the first or second antibody comprises an amino acid
sequence according to Figure 23, 24, 25 or 26.
3. A receptor as claimed in claim 1 or 2 further
comprising:
 - 25 (e) a V_H and C_{H1} region of a third antibody, wherein
the binding specificity of the third antibody
is different from the antigen binding
specificity of the first and second antibody;
 - (f) a polypeptide spacer that links the third
antibody to the first or second antibody; and
 - 30 (g) the light chain of the third antibody.

- 30 -

4. A receptor as claimed in claim 3, wherein binding specificity of the third antibody is directed against DTPA or EDTA.
- 5 5. A process for the preparation of receptors as claimed in claim 1, 2, 3 or 4 comprising the steps of:
 - (a) preparing an expression vector encoding a heavy chain portion of a receptor of claims 1 to 3;
 - (b) preparing an expression vector encoding the respective light chains of a receptor of claims 10 1 to 3;
 - (c) transforming cells with the expression vector from (a) and the expression vector of (b); and
 - (d) growing cells from (c) and isolating the receptor from the cells.
- 15 6. A use of a receptor as claimed in any one of claims 1 to 4 as a pharmaceutical.

FIG. 1






FIG. 2




By: Angus, Buerkin & Parr



FIG. 3





By: Rogers, Brushkin & Parr



By: Rogers, Bureskin & Parr



By: Rogers, Brewster & Parr



By: Rogers, Brushkin & Pan

2019559





FIG.9


By: Rogus, Brestkin & Parr

FIG. 10




By: Rogers, Bruckin & Pan

FIG. 11




By: Rogers, Buehler & Pan

FIG. 12




By: Rogers, Boushkin & Parr

FIG. 13



By: Rigus, Bruehlin & Parr


FIG.14



By: Roger, Berishin & Pass

2019559


FIG. 15



By: Rogers, Bruckin & Pan


2019559

FIG. 16




By: Roger, Brushkin & Parr

FIG. 17




By: Rogers, Bereshkin & Parr

FIG. 18




By: Roger Boushkin & Pam

FIG. 19



By: Rogers, Brashin & Parr


FIG. 20



By: Rogers, Bruckin & Parr

2019559

FIG. 21



By: Roger, Burskin & Parr

FIGURE 22

Peptide linkers with relevant nucleotide sequences

		E	P	K	S	C	G	G	E	A	A	P	A
+	5'	CTCTCTGCAGAGCCCAAATCTTGTGGCGGCGAGGCAGCTCCCGCAG											
-	3'	GAGAGACGTCTCGGGTTTAGAACACCGCCGCTCCGTCGAGGGCGTC											

		A	A	P	A	A	A	A	G	G	Q	V	Q	L	Q	E	S
		CTGCACCCGCAGCAGCCGCAGCAGGCGGGCAGGTCCAAGTGCAGGAGAGC															
		GACGTGGGCGTCGTCGGCGTCGTCCGCCCCTCCAGGTTGACGTCCTCTCG															
																	5'

25

+ = coding strand

- = complementary strand

FIGURE 23

Mab A

VH

Q V Q L Q E S G G G L V Q P G G S L R L
 CAGGTCCAAGTGCAGGAGTCTGGAGGAGGCTTGGTACAGCCTGGGGGTCTCTGAGACTC
 10 20 30 40 50 60

S C A T S G F S D Y Y M N W V R Q P P G
 TCCTGCGCAACTTCTGGGTTCAGTGATTACTACATGAAGTGGGTCCGCCAGCCTCCAGGA
 70 80 90 100 110 120

K A L E W L G F I S N K P N G H T T E Y
 AAAGCACTTGAGTGGTTGGGTTTTATTTCAAACAAACCTAATGGTCACACAACAGAGTAC
 130 140 150 160 170 180

S A S V K G R F T I S R D N S Q S I L Y
 AGTGCATCTGTGAAGGTCGTTTACCATCTCCAGAGATAATTCCCAAAGCATCCTCTAT
 190 200 210 220 230 240

L Q M N T L R A E D S A T Y Y C A R D K
 CTTCAAATGAACACCCTGAGAGCTGAGGACAGTGCCACTTATTATTGTGCAAGAGATAAG
 250 260 270 280 290 300

G I R W Y F D V W G Q G T T V T V S S
 GGAATACGATGGTACTTCGATGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCA
 310 320 330 340 350

VK

A I L S A S P G E K V T M T C R A S S S
 AGCAATCCTGTCTGCATCTCCAGGGGAGAAGGTCACAATGACTTGCAGGGCCAGCTCAAG
 10 20 30 40 50 60

V S Y M H W Y Q Q K P G S S P K P W I Y
 TGTAAGTTACATGCACTGGTACCAGCAGAAGCCAGGATCCTCCCCAAACCCTGGATTIA
 70 80 90 100 110 120

A T S N L A S G V P A R F S G S G S G T
 TGCCACATCCAACCTGGCTTCTGGAGTCCCTGCTCGCTTCAGTGGCAGTGGGTCTGGGAC
 130 140 150 160 170 180

S Y S L T I I R V E A E D A A T Y Y C Q
 CTCTTACTCTCTCAATCATCAGAGTGGAGGCTGAAGATGCTGCCACTTATTACTGCCA
 190 200 210 220 230 240

Q W S S N P L T F G A G T K L E I
 GCAGTGGAGTAGTAACCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGATC
 250 260 270 280 290

FIGURE 24

Mab B

-VH

L Q E S G P D L V K P S Q S L S L T C T
 CTGCAGGAGTCAGGACCTGACCTGGTGAACCTTCTCAGTCACTTTCACCTCACCTGCACT
 10 20 30 40 50 60

V T G Y S I T S G Y S W H W I R Q F P G
 GTCAGTGGCTACTCCATCACCAGTGGTTATAGCTGGCACTGGATCCGGCAGTTTCCAGGA
 70 80 90 100 110 120

N K L E W M G Y I Q Y S G I T N Y N P S
 AACAACTGGAATGGATGGGCTACATACAGTACAGTGGTATCACTAACTACAACCCCTCT
 130 140 150 160 170 180

L K S R I S I T R D T S K N Q F F L Q L
 CTCAAAAGTCGAATCTCTATCACTCGAGACACATCCAAGAACCAGTTCTTCCTGCAGTTG
 190 200 210 220 230 240

N S V T T E D T A T Y Y C A R E D Y D Y
 AATTCAGTGAAGTACTGAGGACACAGCCACATATTACTGTGCAAGAGAAGACTATGATTAC
 250 260 270 280 290 300

H W Y F D V W G A G T T V T V S S
 CACTGGTACTTTCGATGTCTGGGGCGCAGGGACCACGGTCACCGTCTCCTCA
 310 320 330 340 350

VK

L T Q S P A I M S A S L G E E I T L T C
 CTGACCCAGTCTCCAGCAATCATGTCTGCATCTCTAGGGGAGGAGATCACCTAACCTGC
 10 20 30 40 50 60

S T S S S V S Y M H W Y Q Q K S G T S P
 AGTACCAGCTCGAGTGTAAGTTACATGCACTGGTACCAGCAGAAGTCAGGCACTTCTCC
 70 80 90 100 110 120

K L L I Y S T S N L A S G V P S R F S G
 AAACCTCTTGATTATAGCACATCCAACCTGGCTTCTGGAGTCCCTTCTCGCTTCAGTGGC
 130 140 150 160 170 180

S G S G T F Y S L T I S S V E A E D A A
 AGTGGGTCTGGGACCTTTTATTCTCTCAATCAGCAGTGTGGAGGCTGAAGATGCTGGC
 190 200 210 220 230 240

D Y Y C H Q W S S Y P T F G G G T K L E
 GATTATTACTGCCATCAGTGGAGTAGTTATCCACGTTTCGGAGGGGGGACCAAGCTGGAG
 250 260 270 280 290 300

FIGURE 25

Mab C

VH

Q V Q L Q Q S G P E L V K P G A S V K M
 CAGGTCCAAGTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGGCTTCAGTGAAGATG
 10 20 30 40 50 60

S C K A S G Y T F T Y Y V I H W V K Q K
 TCCTGCAAGGCTTCTGGATACACATTCACTTACTATGTTATTCACTGGGTGAAACAGAAG
 70 80 90 100 110 120

P G Q G L E W I G Y I H P Y N A G T E Y
 CCTGGGCAGGGCCTTGAGTGGATTGGATACATTCACTTACAATGCTGGTACTGAGTAC
 130 140 150 160 170 180

N E K F K G K A T L T S D K S S S T A Y
 AATGAGAAGTTCAAAGGCAAGGCCACACTGACTTCAGACAAATCCTCCAGCACAGCCTAC
 190 200 210 220 230 240

M E L S S L T S E D S A V Y Y C S M G R
 ATGGAGCTCAGCAGCCTGACCTCTGAGGACTCTGCGGTCTATTACTGTTCAATGGGACGA
 250 260 270 280 290 300

G G D Y W G Q G T T V T V S S
 GGGGGTGACTACTGGGGCCAAGGGACCACGGTCACCGTCTCCTCA
 310 320 330 340

VK

L T Q S P A I M S A S P G E K V T M T C
 CTGACCCAGTCTCCAGCAATTATGTCTGCATCTCCTGGGGAGAAGGTCACCATGACCTGC
 10 20 30 40 50 60

S A S S S V S Y M H W Y Q Q K S G T S P
 AGTGCCAGCTCAAGTGTAAAGTTACATGCACTGGTACCAGCAGAAGTCAGGCACCTCCCCC
 70 80 90 100 110 120

K R W I Y D T S K L A S G V P A R F S G
 AAAAGATGGATTTATGACACATCCAACTGGCTTCTGGAGTCCCTGCTCGCTTCAGTGGC
 130 140 150 160 170 180

S G S G T S Y S L T I S S M E A E D A A
 AGTGGGTCTGGGACCTCTTACTCTCTCACAATCAGCAGCATGGAGGCTGAAGATGCTGCC
 190 200 210 220 230 240

T Y Y C Q Q W S S N P F T F G A G T K L
 ACTTATTACTGCCAGCAGTGGAGTAGTAACCCATTCACTTCGGCGCGGGGACCAAGCTG
 250 260 270 280 290 300

FIGURE 26

Mab D

VH

A E S G P G L V R L T S L S I T C T V S
 GCAGAGTCAGGGCCTGGCCTGGTGGCCTCACGAGCCTGTCCATCACTTGCACTGTCTCT
 10 20 30 40 50 60

G F S L I S Y G V H W V R Q P P G K G L
 GGCTTTTCATTAATTAGTTATGGTGTACACTGGGTTCCGCCAGCCTCCAGGAAAGGGTCTG
 70 80 90 100 110 120

E W L G V I W A G G S T N Y N S A L M S
 GAGTGGCTGGGAGTAATATGGGCAGGTGGAAGCACAAATTATAATTCCGGCTCTCATGTCC
 130 140 150 160 170 180

R L S I S K D N S K S Q V F L K M N S L
 AGACTGAGCATCAGCAAAGACAACCTCCAAGAGCCAAGTTTTCTTAAAAATGAACAGTCTG
 190 200 210 220 230 240

Q T G D T A I Y Y C A R G G D D Y D G F
 CAACTGGTGACACAGCCATATACTACTGTGCCAGAGGGGGGGATGATTACGATGGGTTT
 250 260 270 280 290 300

A Y W G Q G T T V T V S S G E S
 GCTTACTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGGTGAGTCC
 310 320 330 340

VK

L T Q S P S S L A V S A G E K V T M S C
 CTGACCCAGTCTCCATCCTCCCTGGCTGTGTGCAGCAGGAGAGAAGGTCACTATGAGCTGC
 10 20 30 40 50 60

K S S Q S L L S S T K R K N Y L A W Y Q
 AAATCCAGTCAGAGTCTGCTCAGCAGTACAAAGCGAAGAACTACTTGGCTTGGTACCAG
 70 80 90 100 110 120

Q K P G Q S P K L L I Y W A S T R E S G
 CAGAAACCAGGTCACTCTCCTAACTACTGATCTACTGGGCATCCACTCGGAATCTGGG
 130 140 150 160 170 180

Y P D R F T G S G S G T D F T L T I S S
 GTCCCTGATCGCTTCACAGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGT
 190 200 210 220 230 240

V Q A E D L A V Y Y C K Q S Y N L R A F
 GTGCAGGCTGAAGACCTGGCAGTTTATTACTGCAAACAATCTTATAATCTTCGGGCGTTTC
 250 260 270 280 290 300

G G G T K L E I K
 CGTGGAGGGACCAAGCTGGAGATCAAA
 310 320

FIGURE 27

Mutagenic oligonucleotides:

- 1) 5'CTTACCTGGG.CATGCCCCGA.GCTCCCGTGG.GCATGT3'
- 2) 5'AGTGGGGTTT.TCAGCTCTGCAGAG3'

FIGURE 28

Mutated hinge exon:

^E/_L/^K/_T/^P/_L/^G/_D/^T/_T/^H/_T/^C
 AGAGCTCAAA.ACCCCACTTG.GTGACACAAC.TCACACATGC
 ↓
 G

^P/_R/^C/_P
 CCACGGTGCCCAGGT
 ┌──────────┐
 GAGCTCGGGGCA
 Sst1 Sph1

FIGURE 29

Jun I oligonucleotide

5'CTACGCTCGG.CTAGAGGAAA.AAGTGAAAAC.
CTTGAAAGCG.CAAAACCTCCG.AGCTGGCATC.
CACGGCCAAC.ATGCTCAGGG.AACAGGTGGC.
ACAGCTTAAG.CAGAAAGTCA.TGAACCACCG.
ACCTGCATB3'.

Jun II oligonucleotide

5'CAGGTCGGTG.GTTCATGACT.TTCTGCTTAA.
GCTGTGCCAC.CTGTTCCCTG.AGCATGTTGG.
CCGTGGATGC.CAGCTCGGAG.TTTTGCGCTT.
TCAAGGTTTT.CACTTTTCC.TCTAGCCGAG.
CGATGAGCT3'.